Atrichum altecristatum, 219+; angustatum, 219+; undulatum, 219+
Aulacomniales comb. et stat. nov., 554
Aulacomnium androgynum, 537
Aulaxina microphana, 632+*; quadrangula, 633+*m
Austin protocol (desiccation-tolerance), 171+
Austria (*Bucklandiella nivalis*), 92+
β-orcinol chlorodepsidones (*Lepraria*), 117
Bacidia lobarica sp. nov., 487+*m
Bacidiaceae (*Bacidia*), 487+
Backbone hornwort phylogeny, 188*
Backbone liverwort phylogeny, 186*
Bačkor, Martin, Peter Váczi, Miloš Barták, Jana Buďová and Alexander Dzubai. Uptake, photosynthetic characteristics and membrane lipid peroxidation levels in the lichen photobiont *Trebouxia erici* exposed to copper and cadmium, 100+
Balantiopsis, 746+
Barbella cubensis, *42*; enervis, 42; perpinnata, 41
Barbellopsis trichophora, 42
Bark texture, 770+
Barreno, E. *See* Fernández-Salegui, A. B., 723+
Barták, Miloš. *See* Bačkor, Martin, 100+
Bartholomew-Began, Sharon. *See* Jones, Nicole L., 453+
Bartramia bellolioella sp. nov., 507+*; halleriana, 537; stricta, 537
Bartramiaceae (*Bartramia*), 507+
Base saturation (*Philonotis*), 776+
Baseline physiological responses (*Bazzania*), 297+
Bazzania, 746+; trilobata, 295+
Bednarek-Ochyra, Halina. A taxonomic monograph of the moss genus *Codriophorus* P. Beauv. (Grimmiaceae) [review], 130+
Bednarek-Ochyra, Halina. *See* Köckinger, Heribert, 92+
Bell, Neil E., Dietmar Quandt, Terry J. O'Brien and Angela E. Newton. Taxonomy and phylogeny in the earliest diverging pleurocarps: square holes and bifurcating pegs, 533+
Bell, Neil E. and Peter V. York. *Vetiplanaxis pyrrhobryoides*, a new fossil moss genus and species from Middle Cretaceous Burmese amber, 514+
Bescherellia cryphaeoides, 537; elegantissima, 537
Biodiversity (as air pollution indicator), 724+; (bryophytes in Nevada), 673+; (lichen communities), 724+
Bioindicator species, 727+
Biomass (*Bazzania*), 302+; (lichen litter), 266+
Biomonitoring, 723+; study area species list, 728+
Biotic soil crusts (member taxa), 710+; (site variation), 706+; (species richness), 706+
Blasia, 746+
Blasiales (*Cavicularia*), 453+
Blastenia moreliiensis, 806
Boreal forest (lichens), 74+
Borrera crinalis, *790*
Brachymenium nepalense, 537
Braggins, John E. *See* Hendry, Tory A., 738+
Braithwaitea sulcata, 537
Braithwaiteaceae fam. nov., 554
Branching response (*Bazzania*), 305+
Braun-Blanquet scale, 778
Brinda, John C., Lloyd R. Stark, James R. Shevock and John R. Spence. An annotated checklist of the bryophytes of Nevada, with notes on collecting history in the state, 673+
Bromus tectorum (effect on biotic soil crusts), 708+
Brunet, Jörg. *See* Caldiz, Mayra S., 266+
Bryhnia graminicolor, 759
Bryocaulon pseudosatoanum, 529
Bryophyte genera of Nevada, 704+
Bryophyte papers with phylogeny in the title, 184*
Bryophyte phylogeny, 179+
Bryophytes (biotic soil crusts), 714; (desiccation-tolerance), 595+; (Nevada), 673+
Bryopteris filicina, 766+*
Bryoria spp., 529
Bryum alpinum, 537; argenteum, 537; dendroides, *555*
Buck, William R., Bruce Allen and Ronald A. Pursell. Recent literature on bryophytes—110(1), 133+; 110(2), 337+; 110(3), 567+; 110(4), 838+
Buck, William R. *See* Lücking, Robert, 622+
Bucklandiella, 97+; **sect. Gemmiferae sect. nov.**, 97; **nivalis sp. nov.**, 92+*; vulcanicola (gemmae), 96*
Buďova, Jana. *See* Bačkor, Martin, 100+
Bulgaria (*Philonotis*), 778
Bullatina aspidota, *633*
Buying shed bag sampling (moss), 754+
Cadmium stress (*Trebouxia erici*), 101+
Calciphiles (biotic soil crusts), 714
Caldiz, Mayra S., Jörg Brunet and Bengt Nihlgård. Lichen litter decomposition in *Nothofagus* forest of northern Patagonia: biomass and chemical changes over time, 266+
Calenia aspidota, 633*; solorinoides, 633*
Caliche exposure (biotic soil crusts), 714
California (*Lepraria*), 115+; (lichens and bryophytes), 420+; (*Pleuridium*), 510+
Calomnion brownseyi, 537; complanatum, 537
Caloplaca in North and Central America (key), 799
Caloplaca arenaria, 79+; borealis/kamczatica group, 799; catalinae, 804; cateiloides, 801; cerina, 798+*m; crocea, 799+*m; crocea subsp. mexicana, 802; erythrantha, 799*; ferruginea, 804; haematites, 806; pinicola, 799m; rosulans, *411*; sp. 1, 799*; sp. 2, 799*; stanfordensis, 804; stillicidiorum, 800; xanthostigmoidea, 79+
Calycin (*Candelariella*), 366+, 376+, 392+
Calycularia, 746+
Calymperaceae (evolution), 58; (phylogeny), 46+; (rank-free treatment), 71+
Calymperes, 48+; afzelii, 50; couguiense, 50; dozyanum, 50; erosum, 50; hispidum, 50; lonchophyllum, 50; motleyi, 50; palisotii, 50; robinsonii, 50; schmidtii, 50; tahitense, 50; tenerum, front cover 110(1)*, 50; tuamotuense, 50
Calypogeia, 746+
Candelaria arctica, *388*
Candelariaceae, 365+, 375+, 391+
Candelariella, 365+, 375+, 391+; **aggregata sp. nov.**, 393+*m; antennaria, 396+*m; arctica, 388; athallina, 388+; aurella, 398+*m; aurella group, 391+; **biatorina sp. nov.**, 366+*m; **borealis sp. nov.**, 377+*m; brouardii, 416; **californica sp. nov.**, 400+*m; canadensis, 416+; citrina, 401+*m; **complanata sp. nov.**, 404+*m; coralliza, 389; coralliza f. dispersa, *417*; **corallizoides sp. nov.**, 405+*m; deflexa, *399*; **deppeanae sp. nov.**, 406+*m; dispersa, 417; efflorescens, 379+*m; **immarginata sp. nov.**, 408+*m; kansuensis, 410+*m; kansuensis f. frustulenta, *410*; kuusamoënsis var. areolata, 389; lavicola, 417; **lichenicola sp. nov.**, front cover 110(3)*, 368+*m; litoralis, *399*; lutella, 381+*m; placodizans, 382+*m; plumbea, 417; reflexa, 417; rosulans, 411+*m; rosulans group, 391+; septentrionalis, *383*; spraguei, 415+*m; stenospora, 417; subdeflexa, 365+*m; terrigena, *401*; terrigena var. placodimorpha, *401*; vitellina, 384+*m, 735+; vitellina group, 375+*m; vitellina var. rosulans, *411*; xanthostigma, 386+*m
Canfield line method, 770+*
Cano, María J. *See* Jiménez, Juan A., 812+
Carbon balance (desiccation-tolerance), 605
Cargill, D. Christine. *See* R. Joel Duff, 214+
Carlsson, Bengt Å. *See* Sonesson, Mats, 244+
Cavernularia hultenii, 529; lophyrea, 529
Cavicularia densa, 453+

Cell water relations (bryophytes), 599
Cellular protection (desiccation-tolcrance), 605+
Central America (*Caloplaca*), 798+
Cephalozia, 746+
Cephaloziella, 746+
Ceratolejeunea, 465+*, 746+; dentato-cornuta, *470*; dussiana, *466*
Cetraria californica, 529; chlorophylla, 529
Chaetophyllopsidaceae, 740
Chaetophyllopsis, 746+
Chara vulgaris, 219+
Charales, 612
Cheatgrass (effect on biotic soil crusts), 706+
Chemical changes (lichen), 266+
Chile (*Bartramia*), 507+
Chile (*Didymodon coquimbensis*), 812
Chlorophyll (*Trebouxia erici*), 102
Chlorophyll fluorescence (bryophytes), 163+
Chloroplast genes (liverworts), 738+
Chloroplast sequence markers (*Goniomitrium*), 108+
Chloroplast ultrastructure (phylogeny), 198*+
Cinclidium stygium, 537
Circular tree showing sperm cell evolution, 195*+
Clade A (liverworts), 746+
Clade B (liverworts), 746+
Clade C (liverworts), 746+
Clades (hornworts), 228
Cladina rangiferina, 529
Cladonia spp., 529
Clavitheca nom. nov., 113; **poeltii comb. nov.**, 113
Cleavitt, Natalie L., Karen L. Klima and Timothy J. Fahey. The role of water availability in seasonal growth of *Bazzania trilobata* on boulders in an eastern hemlock stand, 295+
Cleavitt, Natalie L. *See* Proctor, Michael C. F., 595+
Clerc, Philippe and Philip F. May. *Usnea flammea* (Lecanorales) new for North America, 126+
Climate (Tok, Alaska), 77*
Climate effects (lichens and bryophytes), 436+
CLUSTALX, 109, 744
Coeleochaetales, 612
Columbia Basin (biotic soil crusts), 706+
Combined-evidence cladistic analysis, 49+
Commercial moss harvest, 752+
Community analysis (biotic soil crusts), 713+
Community composition (soil-dwelling lichen and moss), 707+
Community dynamics (bryophytes), 760
Community gradients (lichens), 527+
Concentration of elements (lichens), 271+
Connorstictic acid (*Heterodermia*), 492
Conocephalum, 746+
Conservation (pleurocarpous mosses), 327+
Constictic acid (*Usnea flammea*), 126
Copper stress (*Trebouxia erici*), 101+
Cordeiro, V. A., 119
Correspondence analysis, 779
Corynotheca, *113*; poeltii, *113*
Costa (*Orthostichella*), 3+
Costa Rica (*Nothoceros*), 279+
Crust integrity, 715
Crustose lichens (biotic soil crusts), 714
Cryptogams (biomonitoring), 723+
Crytopodium bartramioides, 537
Cultures (*Trebouxia erici*), 101
Cyanolichens (biotic soil crusts), 714
Cyrtomnium hymenophyllum, 537
Cyrtopodendron vieillardii, 537
Cyrtopus setosus, 537
Davis, E. Christine. *See* Hendry, Tory A., 738+
DBH of tree, 771+
DeBolt, Ann M., Roger Rosentreter and Erin P. Martin. Macrolichen diversity in subtropical forests of north-central Florida, 254+
Decay indices, 53
Decomposition parameters (lichens), 268
Degelia plumbea (distribution), 127
Dehydration (bryophytes), 163+
Dehydrins (bryophytes), 606
Dendroceros aff. canaliculatus, 217+; crispatus, 217*+; crispus, 217+; granulatus, 217+; indet., 217+
Dendrocerotaceae (*Nothoceros*), 279+
Dendrogram (lichens and bryophytes), 432+
Dendro-hypnum, 555; **auricomum comb. nov.**, 555; **auricomum ssp. celebensis comb. nov.**, 555+; **auricomum ssp. sumatranum comb. nov.**, 556; beccarii, 556; **brevipes comb. nov.**, 556; **flagelliferum comb. nov.**, 556; **fuscomucronatum comb. nov.**, 556; **fuscomucronatum ssp. chalmersii comb. nov.**, 556; **milnei comb. nov.**, 556; **milnei ssp. korthalsii comb. nov.**, 556; **milnei ssp. parvum comb. nov.**, 556; **opacum comb. nov.**, 556; **reinwardtii comb. nov.**, 556; **reinwardtii ssp. caducifolium comb. nov.**, 556; **subspininervium comb. nov.**, 556; **subspininervium ssp. arborescens comb. nov.**, 556
Derr, Chiska C., Bruce McCune and Linda H. Geiser. Epiphytic macrolichen communities in *Pinus contorta* peatlands in southeastern Alaska, 521+
Desiccation recovery (bryophytes), 600+
Desiccation-tolerance (bryophytes), 595+
Diahyphae (Gomphillaceae), 624*
Dicranum scoparium, 537
Didymodon australasiae, 815; **coquimbensis sp. nov.**, 812+*; fuscus, 815; luehmannii, 815; nevadensis, 815
Diplophyllum, 746+
Disjunction (*Usneae* distribution), 127
Dispersal (desiccation-tolerance), 608+
Distribution (*Bucklandiella*), 99; (lichens of Tetlin National Wildlife Refuge), 80; (*Usneae*), 127
Disturbance (biotic soil crusts), 715+
Ditrichaceae (*Pleuridium*), 510+
Diversity (boreal zone lichens), 89; (desiccation-tolerance), 598
DNA extraction and amplification (hornworts), 219+
DNA sequencing and alignments (Calymperaceae), 51
Drought-hardening, 602+
Drying rate (desiccation-tolerance), 602+
Duckett, Jeffrey G. *See* Renzaglia, Karen S., 179+
Duff, R. Joel, Juan Carlos Villarreal, D. Christine Cargill and Karen S. Renzaglia. Progress and challenges toward developing a phylogeny and classification of the hornworts, 214+
Duff, R. Joel. *See* Renzaglia, Karen S., 179+
Dzubai, Alexander. *See* Bačkor, Martin, 100+
Eastern North America (Gomphillaceae), 622+
Echinoplaca areolata sp. nov., front cover 110(4)*, 633+*m; furcata subsp. neotropica, 635*; intercedens, 635; lucernifera, 635*; pellicula, 635*; similis, 635, 637*; tetrapla, 637*
Ecological niche modelling (*Philonotis*), 777+
Ecophysiology (bryophytes), 599
Egan, Robert S. Recent literature on lichens—204, 146+; —205, 351+; —206, 577+; —207, 824+
Elix, John A. *See* Knudsen, Kerry, 115+
Embryophytes (desiccation-tolerance), 612+
Endocarpon pusillum, 79+
Environmental measurements (*Bazzania*), 297+
Environmental pollution (lichen), 100+
Ephemerum sessile, 109+
Epilithic bryophyte communities, 753+
Epiphyte communities (lichens and bryophytes), 420+
Epiphytic lichens (biomonitoring), 724+
Epiphytic liverwort (spatial distribution), 767+

Epixylic bryophyte communities, 753+
Equihua, Clementina and Miguel Equihua. Spatial distribution of *Bryopteris filicina* on tree trunks in the Selva Lacandona, Chiapas, Mexico, 766+
Equihua, Miguel. *See* Equihua, Clementina, 766+
Erythrodontium subbiformis, 42
Esslinger, Theodore L. A synopsis of the North American species of *Anaptychia* (Physciaceae), 788+
Establishment (desiccation-tolerance), 608+
Europe (*Philonotis*), 776+
Evernia prunastri, 733+
Evolution (desiccation-tolerance), 611+; (liverworts), 748
Excipular types (Gomphillaceae), 624*
Exemplar approach (morphology), 51
Exodictyon incrassatum, 50
Exostratum blumii, 50
Fahey, Timothy J. *See* Cleavitt, Natalie L., 295+
Fernández-Salegui, A. B., A. Terrón, E. Barreno and P. L. Nimis. Biomonitoring with cryptogams near the power station of La Robla (León, Spain), 723+
Field collections (boreal lichens), 76+
Field moss sampling, 755+
Fire (effect on biotic soil crusts), 706+
Fisher, Kirsten M., Dennis P. Wall, Kwok Leung Yip and Brent D. Mishler. Phylogeny of the Calymperaceae with a rank-free systematic treatment, 46+
Florida (lichens), 254+
Folioceros fuciformis, 218*+
Forests, old-growth (*Neckera*), 309+
Fossil moss (*Vetiplanaxis*), 518+
Fossombronia from Latin America (typifications), 817+
Fossombronia, 746+; brasiliensis, 818; carinata, 819; crassifolia, 819; cubana, 819; fernandeziensis, 819; foveolata, 218+; herzogii, 819; lophoclada, 819; luetzelburgiana, 819; mexicana, 819; ptychophylla, 819; tenuifolia, 819; wrightii, 819
Fossombroniaceae, 817+
Freire, A. Virginia and Raymond E. Stotler. Typifications of the names of some Latin American species of *Fossombronia* (Fossombroniaceae), 817+
Frisch, Andreas, Klaus Kalb and Martin Grube. 2006. Contributions towards a new systematics of the lichen family Thelotrematáceae [review], 332+
Frullania, 746+
Funaria hygrometrica, 537
Funariaceae (phylogeny), 110+
Funarioideae, 112+
Future challenges (phylogeny), 205+
Gackstroemia, 746+
Gametangium development (desiccation-tolerance), 608+
Gametophyte development (*Cavicularia*), 453+
Gametophyte evolution, 181+
Garovaglia elegans, 537
Geiser, Linda. *See* Derr, Chiska C., 521+
Gemma/gemmaling ontogeny (*Cavicularia*), 453+*
Gemmae (*Bucklandiella nivalis*), 94+*
Gene expression (desiccation-tolerance), 608+
Gene ontology mapping (bryophytes), 608+
Gene partition congruence and combining data (hornworts), 220
General classification (hornworts), 238+
Geocalyx, 746+
Global diversity (pleurocarpous mosses), 319+
Gloeoheppia squamulosa comb. nov., 291+*
Glyphothecium sciuroides, 537
Goebeliella, 746+
Goebelobryum, 740; unguiculatum, 738+
Goffinet, Bernard. *See* Werner, Olaf, 108+
Gomphillaceae, 622+; (*Gomphillus*), 475+
Gomphillaceae in North America (checklist), 626+; (key), 625+
Gomphillus, 622+; americanus, front cover 110(4)*, 637*m; calycioides, 638*m; **hyalinus comb. nov.**, 476+; ophiosporus, *475+*
Goniobryum subbasilare, 537
Goniomitrium, 108+; acuminatum, 109+; seroi, 109+
Goward, Trevor and Bruce McCune. *Hypogymnia canadensis* (Parmeliaceae), a new lichen from the Pacific Coast of North America, 808+
Graphids, Australian (lichens), 561+
Greven, Henk. *See* Hastings, Roxanne I., 500+
Grimmia afroincurva, 500+; donniana, 504; incurva, 504; **milleri sp. nov.**, 500+*; trichophylla, 93
Grimmiaceae (gemmae), 96; (*Grimmia*), 500+
Groundwater pH, 778
Growth (*Neckera*), 309+
Gyalectidium, 623+; appendiculatum, 639*; aspidotum, *633*; catenulatum, 640*; denticulatum, 642; eskuchei, 642; filicinum, 641*; floridense, 641*; imperfectum, 641*; kenyanum, 642; **tuckerae sp. nov.**, front cover 110(4)*, 641*; **viride sp. nov.**, 643+*; yahriae, 643*
Gyalideopsis, 622+; sp. A, 650+*; sp. B, 663; anastomosans, *663*; aequatoriana, 657; africana, 643; alnicola, 656; **americana sp. nov.**, 643+*m; aff. argentea, 645*; **berenice comb. nov.**, 645*; buckii, 647*m; **epicortis comb. nov.**, 650*; floridae, 650*m; gigantea, 647; glauca, 655; lambinonii, 651*m; macarthurii, 653*; moodyae, 653*m; muscicola, 653; **ozarkensis sp. nov.**, front cover 110(4)*, 655+*m; piceicola, 656; **subaequatoriana sp. nov.**, 657+*m; **submonospora sp. nov.**, front cover 110(4)*, 659+*m; aff. vainioi, 663; **vainioi var. semicirculata var. nov.**, 652+*m; vainioi var. vainioi, 652+*m; wesselsii, 657+*
Gyalolechia brouardii, *416*
Gyrothyra, 746+
Habitat differentiation (*Philonotis*), 776+
Habitat structure (lichens and bryophytes), 428+
Hájek, Michal. *See* Hájková, Petra, 776+
Hájková, Petra, Blanka Shaw, Michal Hájek, Dirk Hinterlang and Vítězslav Plášek. The role of base saturation and altitude in habitat differentiation within *Philonotis* in springs and mires of three different European regions, 776+
Hampeella alaris, 537
Haplomitrium (phylogeny), 192*+
Haplomitrium hookeri, 746+; mnioides, 746+
Hardwood bottom vegetation (lichens), 263+
Harmand, Julien, 119
Harpanthus, 746+
Harris, Richard C. *See* Lendemer, James C., 490+
Harvest methods (moss), 753
Hassel de Menéndez, Gabriela. *See* Villarreal, Juan Carlos, 279+
Hastings, Roxanne I. and Henk Greven. *Grimmia milleri* sp. nov. (Grimmiaceae) from northeastern North America and the status of *Grimmia afroincurva*, 500+
Hattorianthus, 746+
Heavy metals (lichen), 100+
Hedenäs, Lars. Global diversity patterns among pleurocarpous mosses, 319+
Hedwigia ciliata, 537
Hemlock stand (*Bazzania*), 295+
Hendry, Tory A., Bin Wang, Ya Yang, E. Christine Davis, John E. Braggins, Rudolf M. Schuster and Yin-Long Qiu. Evaluating phylogenetic positions of four liverworts from New Zealand, *Neogrollea notabilis*, *Jackiella curvata*, *Goebelobryum unguiculatum* and *Herzogianthus vaginatus*, using three chloroplast genes, 738+
Heppia despreaxii, 287+*
Herbarium websites, 676
Herbertus, 746+

Herzogianthus, 746+; vaginatus, 738+
Heterodermia neglecta sp. nov., 490+*
Hildebrandtiella endotrichelloides, 537; pachyclada, *42*; **phleoides comb. nov.**, 41
Hill, David J. 2006. Surveying and report writing for lichenologists [review], 564+
Hinterlang, Dirk. *See* Hajková, Petra, 776+
History of bryophyte collection (Nevada), 673+
Hohe Tauern National Park (*Bucklandiella*), 93
Holien, Håkon and Tor Tønsberg. Norsk lavflora [review], 129+
Homeohydry, 598
Homoiohydry, 598
Hookeria lucens, 537
Hornwort (*Nothoceros*), 279+
Hornworts (desiccation-tolerance), 163+; (phylogeny), 214+
Horse Heaven Hills (Washington), 708+m
Hubbard Brook (*Bazzania*), 295+
Huisman, Olff and Fresco (HOF) type 5 model, 779
Humidity (*Bryopteris filicina*), 772+
Huperzia lucidula, 219+; selago, 219+; verticillata, 219+
Hydrins (desiccation-tolerance), 607
Hygrolejeunea ocellata, *470*
Hymenodon pilifer, 537; sphaerothecius, 537
Hymenodontopsis **bifaria comb. nov.**, 555; **mnioides comb. nov.**, 555; stresemannii, 537; **vallis-gratiae comb. nov.**, 555
Hymenophyton, 746+
Hyphophore development, 622+
Hyphophores, front cover 110(4)*, 622+
Hypnodendrales comb. et stat. nov., 554
Hypnodendron arcuatum, 537; auricomum, 537, *555*; auricomum ssp. celebensis, *556*; auricomum ssp. opacum, *556*; auricomum ssp. sumatranum, *556*; beccarii, *556*; brevipes, *556*; caducifolium, *556*; camptotheca, 537, *555*; chalmersii, *556*; colensoi, *555*; comatulum, 537, *555*; comatum, 537, *555*; comosum, 537, *555*; comosum var. seiberi, *555*; dendroides, 537, *555*; diversifolium, 537, *557*; flagelliferum, *556*; fuscomucronatum, 537, *556*; fuscomucronatum ssp. chalmersii, *556*; kerrii, 537, *556*; menziesii, 537, *556*; microstictum, 537; milnei, 537, *556*; milnei ssp. korthalsii, *556*; milnei ssp. parvum, *556*; opacum, *556*; reinwardtii ssp. caducifolium, *556*; reinwardtii, 537, *556*; spininervium, 537; subspininervium, 538, *556*; subspininervium ssp. arborescens, *556*; sumatranum, *556*; tahiticum, *555*; vitiense, 538
Hypnum bifarium, *555*; comatum, *555*; comosum, *555*; cupressiforme, 538; curvifolium, 756+; divaricatum, *555*; fuscomucronatum, *556*; hexastichum, *2+*; imbricatum, *29*; imponens, 756+; menziesii, *556*; mnioides, *555*; pentastichum, *41*; phleoides, *42*; reinwardtii, *556*; seiberi, *555*; subspininervium, *556*
Hypogymnia apinnata, 529; **canadensis sp. nov.**, 808+*m; duplicata, 529; enteromorpha, 529; imshaugii, 529; inactiva, 529; krogiae, 811; metaphysodes, 808+; oceanica, 529; sachalinensis, 811; vittata, 529
Ilkiu-Borges, Anna Luiza. *See* Reiner-Drehwald, M. Elena, 465+
Illegal harvest (moss), 753
Incidental take (moss harvest), 752+
Indicator species analyses (lichens and bryophytes), 427+
Indicator species analysis, 712, 756
Inductively coupled plasma atomic emission spectrometry, 101
Ingerpuu, Nele, Kai Vellak and Tõnu Mõls. Growth of *Neckera pennata*, an epiphytic moss of old-growth forests, 309+
Intermittent desiccation (bryophytes), 604+
Invertebrates (moss harvest), 761
Invited Essay: New Frontiers in Bryology and Lichenology, 163+
Invited review: Desiccation-tolerance in bryophytes, 595+
Ireland, Robert R. *See* Allen, Bruce, 506+
Isoetes engelmannii, 219+; histrix, 219+; kirkii, 219+
Isophylly, 748
Isotachis, 746+
Isothecium colensoi, *555*
Isousnic acid (*Lecanora*), 485
Jaccard's indices of similarity (pleurocarpous mosses), 320+
Jackiella curvata, 738+
Jackiellaceae, 739+
Jamesiella anastomosans, 663*m
Jensenia, 746+
Jiménez, Juan A. and Maria J. Cano. *Didymodon coquimbensis* (Pottiaceae), a new species from Chile, 812+
Jones, Nicole L. and Sharon Bartholomew-Began. Juvenile gametophyte development in the Blasiales (Metzgeriidae). 2. Gemma/gemmaling ontogeny of *Cavicularia densa*, 453+
Jovan, Sarah. [Review of:] Lambley, Peter and Pat Wolseley. 2004. Lichens in a changing pollution environment, 333+
Jubula, 746+
Julella vitrispora, 120
Jungermannia, 746+
Kellman, Ken M. *See* Yip, Kwok Leung, 510+
Key (families and genera of the Hypnodendrales), 557+; (genera and species of Gomphillaceae in North America), 625+; (*Gloeoheppia* and *Pseudopeltula* in southwestern North America, Central America, and the Caribbean), 293+; (lichen genus *Candelariella* in western North America), 392+; (polysporous *Candelariella* species in Western North America), 376+; (species of *Anaptychia* in North America), 789+; (species of *Caloplaca* in North and Central America), 799; (species of *Orthostichella*), 5; (species of *Pleuridium* in North America), 512+
Klima, Karen L. *See* Cleavitt, Natalie L., 295+
Knudsen, Kerry and John A. Elix. A new *Lepraria* (Stereocaulaceae) from the Santa Monica Mountains in southern California, 115+
Knudsen, Kerry. [Review of:] McCune, B. and R. Rosentreter. 2007. Biotic soil crust lichens of the Columbia Basin, 563+
Köckinger, Heribert, Halina Bednarek-Ochyra, and Ryszard Ochyra. Bucklandiella nivalis (Grimmiaceae), a new moss from the Alps of Austria, 92+
Kondratyuk, Sergey Ya. In memoriam: Maria F. Makarevych. D. Sc., Professor (December 4, 1906–February 24, 1989), 480+
Kriging, 727
Kučera, Jan. [Review of:] Schumacher, René and Jiri Váňa. 2005. Identification keys to the liverworts and hornworts of Europe and Macaronesia (distribution and status), 335+
La Robla, Spain (biomonitoring), 723+m
Lacandona, Mexico (*Bryopteris filicina*), 766+
Lambley, Peter and Pat Wolseley. 2004. Lichens in a changing pollution environment [review], 333+
Landscape strata (biotic soil crusts), 715+
Late Embryogenesis Abundant protein group, 606
Latin America (*Fossombronia*), 817+
Lawton, Elva, 673
LEA protein group (bryophytes), 606
Leafy liverworts (phylogeny), 746+
Lecanora abellula, 483+; citrina var. xanthostigma, *386*; coniferarum, 483+; conjungens, *802*; coralliza, *389*; **crassithallina sp. nov.**, 482+*; deflexa, *399*; erythranta, *802*; lividocinerea, 120; manuelina syn. nov., *120*; russeola, *802*; saligna, 483+; vitellina f. athallina, *388*; vitellina var. placodizans, *382*; vitellina var. reflexa, *417*
Lecanoraceae (*Lecanora*), 482+
Lecanorales (*Bacidia*), 487+
Lecidea crocea, *801*
Lecidella elaeochroma, 733+
Leiomitra, 746+
Leiosporoceros (phylogeny), 189*+; dussii, 218+

Lejeunea, 746+; acanthotis, *470*; huctumalcensis, 465+*; schizotis, *470*; xiphotis, *470*
Lejeuneaceae (*Lejeunea*), 465+
Lembophyllaceae, 3
Lempholemma chalazanum, 289+*
Lendemer, James C., Richard C. Harris and Erin A. Tripp. *Heterodermia neglecta* (Physciaceae), a new lichen species from eastern North America, 490+
Lepicolea, 746+
Lepidogyna, 746+
Lepidolaena, 746+
Lepidozia, 746+
Lepidoziaceae, 739+; (*Meinungeria*), 494+
Lepraria coriensis, 117; **santamonicae sp. nov.**, 115*
Leptostomum inclinans, 538
Leptotheca boliviana, 538; gaudichaudii, 538+*
Leucobryoid taxa, 48+
Leucodon sciuroides, 538
Leucolepsis acanthoneura, 538
Leucophanes glaucum, 50; octoblepharoides, 50; seychellarum, 50
Lichen diversity, 725+
Lichen litter, 266+
Lichen photobiont (photosynthesis), 100+
Lichen stillicidiorum, *799*
Lichen, sorediate (*Bacidia*), 487+
Lichens (Alaska), 74+; (biomonitoring), 724+; (Portugal), 119+
Lichens of Tetlin National Wildlife Refuge and area (checklist), 79+
Ligrone, Roberto. *See* Renzaglia, Karen S., 179+
Linear mixed model analysis (*Bazzania*), 303+
Linnaean system, 47+
List (hornworts with desiccation-tolerance), 173; (liverworts with desiccation-tolerance), 172+; (macrolichen species, north-central Florida), 259+; (mosses with desiccation-tolerance), 166+; (nonbryophyte species affected by moss harvest), 762; (species directly impacted by commercial moss harvest), 757+; (species in biomonitoring study area), 728+; (taxa on *Sequoia sempervirens* in old growth forest), 443+; (bryophytes in Nevada), 677+; (Gomphillaceae in North America), 626+ (lichens of Tetlin National Wildlife Refuge and area), 79+;
Liverworts (desiccation tolerance), 163+, 613*; (phylogenetic analysis), 738+
Lobaric acid (*Bacidia*), 488; (*Usnea flammea*), 126
Local mean models, 712
Log moss (harvest), 753+
Loiseaubryum, 113; ephemeroides, 113
Longleaf pine vegetation (lichens), 262+
López de Silanes, M^a Eugenia. *See* Paz-Bermúdez, Graciela, 119+
Lophocolea, 746+; cuspidata, 759
Lopidium concinnum, 538
Lücking, Robert. The fungi *Microstelium hyalinum* and *Acrospermum puiggarii* are the same as the lichen *Gomphillus ophiosporus* (Ostropales: Gomphillaceae), 475+
Lücking, Robert, William R. Buck and Eimy Rivas Plata. The lichen family Gomphillaceae (Ostropales) in eastern North America, with notes on hyphophore development in *Gomphillus* and *Gyalideopsis*, 622+
Lunularia, 746+
MacClade 4.05, 744
Macrocoma tenuis, 538
Macrolichen communities, epiphytic, 521+
Macrolichen diversity (Florida), 254+
Magill, Robert E. *See* Allen, Bruce, 1+
Maine, coastal (*Usnea flammea*), 126+
Makarevych, Maria F. (memorial), 480+*
Malcolm, B. and N. Malcolm. 2006. Mosses and Other Bryophytes—An Illustrated Glossary. 2nd Edition [review], 566
Mangold, Armin. [Review of:] Frisch, Andreas, Klaus Kalb and Martin Grube. 2006. Contributions towards a new systematics of the lichen family Thelotremataceae, 332+
Marchantia, 746+
Marsupella, 746+
Marsupidium, 746+
Marsupium-bearing clade (liverworts), 746+
Martin, Erin P. *See* DeBolt, Ann M., 254+
Mastigophora, 746+
Maximum likelihood analysis (liverworts), 745+
Maximum parsimony analysis (liverworts), 745+
Maximum parsimony trees (Calymperaceae), 54+*
May, Philip F. *See* Clerc, Philippe, 126+
McCune, B. and R. Rosentreter. 2007. Biotic soil crust lichens of the Columbia Basin. [review], 563+
McCune, Bruce. *See* Derr, Chiska C., 521+; *see* Goward, Trevor, 808+; *see* Ponzetti, Jeanne M., 706+
Megaceros denticulatus, 217+; flagellaris, 217*+; gracilis, 217+; pellucidus, 217+
Meinungeria mouensis gen. et sp. nov., 494+*m
Membrane lipid peroxidation (*Trebouxia erici*), 100+
Memorial: Lewis Edward Anderson, 364
Memorial: Maria F. Makarevych, 480+*
Menegazziaic acid (*Usnea flammea*), 126
Mesochaete taxiforme, 538; undulata, 538
Meteoriaceae, 2+
Meteorium deppei, 41
Metzgeria, 746+; conjugata, 219+, 759
Metzgeriidae (*Cavicularia*), 453+
Mexico (*Bryopteris filicina*), 766+
Mexico, northern (*Candelariella*), 365+, 375+, 391+; (*Lecanora*), 482+
Microclimate (bryophytes), 598
Microenvironmental variables, 770+
Microlychnus epicortis, *650*
Microstelium hyalinum, *475+**
Middle Cretaceous Burmese amber (*Vetiplanaxis*), 518+
Mid-west Europe, 778
Mielichhoferia elongata, 538
Mires (Europe), 776+
Mishler, Brent D. *See* Fisher, Kirsten M., 46+; *see* Proctor, Michael C. F., 595+; *see* Renzaglia, Karen S., 179+
Mitthyridium (monophyly), 56; constrictum, 50; fasciculatum, 50; jungquilianum, 50; obtusifolium, 50; undulatum, 50
Mniodendron, 555; camptotheca, 555; colensoi, 555; comatulum, 555; comatum, 555; comosum, 555; **comosum var. seiberi comb. nov.**, 555; dendroides, 555; korthalsii, 556; parvum, 556; tahiticum, 555
Mnium hornum, 538; vallis-gratiae, 555
Modified CTAB extraction protocol, 744
Moerckia, 746+
Molecular markers (hornworts), 221
Mols, Tõnu. *See* Ingerpuu, Nele, 309+
Monoclea, 746+; gottschei, 219+; indet., 219+
Monodactylopsis, 499
Monongahela National Forest (moss harvest), 753+
Monophyly (Funariaceae), 110+
Morphological reduction (mosses), 112
Morphology (*Orthostichella*), 3+
Moss (Alps of Austria), 92+
Moss harvest (Appalachian Mountains), 752+m
Moss systematics, 47+
Mosses (desiccation-tolerance), 163+, 613*
MrBayes 3.0, 110
Mt. Sauleck (*Bucklandiella nivalis*), 93*
Muller, Frank. *Meinungeria mouensis* (Lepidoziacae), a new genus and species from New Caledonia, 494+

Multiresponse permutation procedure, 712, 756
Myurium hochstetteri, 538
Nanomitriella, 113
Neckera ampullacea, *25*; cyathipoma, *18*; decolorans, *37*; moenkemeyeri, *30*; pachygaster, *30*; pachygaster var. gracilis, *30*; pandurifolia, *21*; pennata, 309+m; rigida, *25*; trichophoroides, *42*; turgidula, *42*; versicolor, *30*; viridis, *30*; welwitschii, *37*
Neckeraceae, 1+
Negrollea notabilis, 738+
Neogrolleoideae, 748
Neotrichocolea, 746+
Neotropics (*Lejeunea*), 465+
Nephroma arcticum, 244+
Nevada (bryoflora), 673+
Nevada (bryophyte checklist), 677+ ; (early bryological exploration), 675m; (index to bryophyte genera), 704+
New Caledonia (*Meinungeria*), 494+
New World (*Orthostichella*), 1+
New Zealand (liverworts), 738+
NEWTON, ANGELA. *See* BELL, NEIL E., 533+
Niche differentiation (*Philonotis*), 784
NIHLGÅRD, BENGT. *See* CALDIZ, MAYRA S., 266+
NIMIS, P. L. *See* FERNÁNDEZ-SALEGUI, A. B., 723+
Nitella flexilis, 219+
NO_2, 724+
Nonbryophyte species affected by moss harvest (list), 762
Nonmetric multidimensional scaling, 712, 756; (pleurocarpous mosses), 320+
Nonparametric multiplicative regression (lichens), 526+
Nontimber forest products, 753+
Norargopsin (*Lepraria*), 117
Norstictic acid (*Heterodermia*), 492; (*Usnea flammea*), 126
North America (*Anaptychia*), 788+; (*Caloplaca*), 798+; (*Gomphillaceae*), 622+; (*Hypogymnia canadensis*), 808; (*Sphagnum*), 274+; (*Sphagnum oregonense*), 122; (*Usnea flammea*), 126+
North America, eastern (*Grimmia*), 500+
North America, eastern (*Heterodermia*), 490+; (*Sphagnum*), 122
Nothoceros cf. aenigmaticus, 217+; fuegiensis, 217+; **canaliculatus comb. nov.**, 283; endiviaefolius, 217*+, **endiviaefolius comb. nov.**, 283; giganteus, 217+, **giganteus comb. nov.**, 283; indet., 217+; **superbus sp. nov.**, 279+*; cf. vincentianus, 217+
Nothofagus forest (lichens), 266+
Notothylas breutelii, 218*+; cf. javanica, 218+; orbicularis, 218+
O'BRIEN, TERRY L. *See* BELL, NEIL E., 533+
Oceanic distribution (*Usneae*), 127
OCHYRA, RYSZARD. *See* KOCKINGER, HERIBERT, 92+
Octoblepharum albidum, 50
Odontoschisma, 746+
OLIVER, MELVIN J. *See* PROCTOR, MICHAEL C. F., 595+
Ontogeny (*Cavicularia*), 453+
Ophioglossum reticulatum, 219+; vulgatum, 219+
Optimality criteria applied to data partitions (hornworts), 220+
Ordination (lichens), 261+
Ordination analysis, 756
Oregon (*Sphagnum oregonense*), 122
Orthodontiales comb. et stat. nov., 554
Orthodontium lineare, 538+*
Orthodontopsis bardunovii, 538
Orthostichella (Key to species), 5
Orthostichella, 1+; **capillicaulis comb. nov.**, 5+*; dusenii, *15*; filamentosula, *31+*; hexasticha, 5+*; leptopteridea, *6*; **longinervis comb. nov.**, 5+*; microcarpa, *26*; mucronatula, *18*; **muelleri comb. nov.**, 5+*; **pachygastrella comb. nov.**, 5+*; pandurifolia, 5+*; **rigida comb. nov.**, 5+*; tijucae, *42*; versicolor, 5+*; **welwitschii comb. nov.**, 5+*
Orthostichidium involutifolium, 25; **pentastichum comb. nov.**, 41+
Orthostichopsis, 13; chrysoneura, 40; tenuis, 40
Osmunda cinnamomea, 219+; regalis, 219+
Ostropales (*Gomphillus*), 475+
Oxidative metabolism (bryophytes), 606
Oxygen evolution rate (*Trebouxia erici*), 102
Pacific Coast (*Hypogymnia canadensis*), 808+
Pallavicinia, 746+
Paraphyses (*Caloplaca*), 798+
Paraschistochila, 746+
Parmelia exasperatula, 733+; hygrophila, 529; saxatilis, 529; spp., 529; squarrosa, 529; sulcata, 529, 733+; tiliacea, 733+
Parmeliaceae, 808+
Parmelion community, 727+
Parmotrema (substrate), 651
Parsimony analysis (pleurocarps), 541+
Particles (air pollution), 724+
Patagonia (lichens), 266+
Patellaria vitellina, *384*
PAUP*4.0b10, 52, 109, 745
PAZ-BERMÚDEZ, GRACIELA AND M^{A} EUGENIA LÓPEZ DE SILANES. The identity of two lichen species described from Portugal by J. Harmand, 119+
PCR, 744
PCR reactions (*Goniomitrium*), 109+
Peatlands (lichens), 521+
PECK, JERILYNN E. *See* STUDLAR, SUSAN MOYLE, 752+
Pellia, 746+
Perennial grasslands (biotic soil crusts), 708+
Petalophyllum, 746+
pH (moss desiccation-tolerance), 599
Phaeoceros carolinianus, 218+; cf. carolinianus, 218+; hallii, 218+; indet., 218+; laevis, 218+; mohrii, 218+; pearsonii, 218+
Phaeomegaceros coriaceus, 217+; fimbriatus, 217+; aff. fimbriatus, 217*+; fimbriatus var. β schlimianus, 217+; hirticalyx, 217+; indet., 217+; skottsbergii, 217+
Phaeophyscia constipata, 790; hirsuta, 733+; orbicularis, 735+
Phaeophytinization quotient (photobiont chlorophyll), 102
Philonotis caespitosa, 777+; calcarea, 777+; fontana, 777+; marchica, 779; seriata, 777+; species (habitat differentiation), 776+; tomentella, 779
Photosynthesis (desiccation-tolerance), 600+; (*Trebouxia erici*), 100+
Phycolepidozia, 499
Phyllodrepanium falcifolium, 538
Phyllothallia, 746+
PhyloCode, 60+
Phylogenetic analyses (Bayesian: hornworts), 221; (liverworts), 744; (parsimony: hornworts), 221; (pleurocarps), 539+
Phylogenetic positions (liverworts), 738+
Phylogenetic reconstructions (hornworts), 224*+
Phylogeny (Calymperaceae), 46+; (pleurocarps), 533+
Phylogeny of land plants (desiccation-tolerance), 613*
Phymatoceros bulbiculosus, 218*+; phymatodes, 218+
PHYML version 2.4.4, 745
Physcia adscendens, 735+; aipolia, 733+; ciliaris var. crinalis, *791*; elbursiana, *792*; grisea var. elbursiana, *792*; ulotrichoides, *794*
Physciaceae, 788+; (*Heterodermia*), 490+
Physcomitrella patens, 538
Physconia enteroxantha, 733+; grisea, 733+; muscigena, 790; petraea, 790; subpallida, 794; thomsonii, *792*
Physiology (*Nephroma*), 244+
Physodic acid (*Hypogymnia canadensis*), 810
Phytogeographic categories (Alaskan lichens), 79+
Pigments (*Heterodermia*), 492; (*Trebouxia erici*), 102
Pilotrichella, 2; allionii, *32*; angustifolia, *32*; attenuata, *37*;

biformis var. congesta, *42*; caldensis, *30*; calomicra, *25*; capillicaulis, *5*; chlorothrix, *42*; chrysoneura, *13*; communis, *31*; conferta, *22*; curvifrons, *5*; cuspidata, *22*; cyathipoma var. laxiretis, *18*; delicatula, *6*; densiramea, *43*; dewrevrei, *37*; eustichica, *41*; gracilicaulis, *31*; grimaldii, *30*; guineensis, *30*; holstii, *26*; illecebraria, *41*; imbricatula var. nervosa, *26*; incurva, *22*; inflatifolia, *30*; isleana var. virescens, *25*; kraussei, *41*; kuntzei, *21*; latiramea, *31*; leptoclada, *25*; lepto-erioclada, 41; longinervis, *10*; mascarenica, 2; mexicana, *25*; muelleri, *15*; obovata, *25*; pachygastrella, *16*; percapillaris, 41; percordata, *6*; perpinnata, *41*; pilifolia, *37*; profusicaulis, *25*; pulchella, *25*; rehmanni, *43*; rigens, *18*; sericea, *26*; serricola, *43*; sordidoviridis, *37*; stuhlmannii, *5*; subimbricata, *13*; subimbricata var. flageyi, *31*; subpachygastrella, *18*; subpachygastrella var. minor, *18*; subpanduraefolia, *26*; tenella, *9*; tenellula, *5*; turgidellacea, *31*
Pinus contorta (lichens), 521+m
Placodium aureofulvum, *802*; brebissonii var. microspora, *806*; cerinum ulmorum, *799*; gilvum var. erythranthoides, *802*; spraguei, *415*
Plagiobryum zierii, 538
Plagiochila, 746+; adiantoides, 219+; asplenoides, 219+
Plagiomnium cuspidatum, 538
Plášek, Vítězslav. *See* Hájková, Petra, 776+
Platismatia glauca, 529; herrei, 529; lacunosa, 529; norvegica, 529
Pleospora vitrispora, *120*
Pleuridium acuminatum, 512; mexicanum, 510+; ravenelii, 512; subulatum, 513; sullivantii var. mexicanum, *510*
Pleurocarpous mosses (global diversity), 319+m
Pleurocarps (taxonomy and phylogeny), 533+
Pleurozia, 746+
Pleurozium schreberi, 538
Pohlia nutans, 538
Poikilohydry, 163+, 598+
Polyblastiopsis cordeiroi syn. nov., *120*
Polymerase chain reaction (Calymperaceae), 51+
Polysporus species (*Candelariella*), 375+
Ponzetti, Jeanne M., Bruce McCune and David A. Pyke. Biotic soil crusts in relation to topography, cheatgrass and fire in the Columbia Basin, Washington, 706+
Porella, 746+
Pores (*Sphagnum oregonense*), 123
Porotrichum, 13
Portugal (lichens), 119+
Pottiaceae, 812; (phylogeny), 111+
Powellia involutifolia, 538
Power station (biomonitoring), 723+
Principal Component Analysis (biomonitoring), 731
Printzen, Christian and Tor Tønsberg. *Bacidia lobarica* (Bacidiaceae, Lecanorales) sp. nov., a sorediate lichen from the southeastern U.S.A., 487+
Proctor, Michael C. F., Melvin J. Oliver, Andrew J. Wood, Peter Alpert, Lloyd R. Stark, Natalie L. Cleavitt and Brent D. Mishler. Desiccation-tolerance in bryophytes: a review, 595+
Protein synthesis (desiccation-tolerance), 607+
Pseudevernia furfuracea, 735+
Psoroma palmulata, *793*
Psorotichia segregata syn. nov., *286+*; squamulosa, *286+*
Pterobryella praenitens, 538
Pterobryon densum, 538
Pteropsiella, 499
Ptilidium, 746+
Ptychomnion cygnisetum, 538
Pulvic acid lactone (*Candelariella*), 366+, 376+, 392+
Pulvinic acid (*Candelariella*), 366+, 376+, 392+
Pursell, Ronald A. [Review of:] Malcolm, B. and N. Malcolm. 2006. Mosses and other bryophytes—An illustrated glossary. 2nd edition, 566
Pursell, Ronald A. *See* Buck, William R., 133+, 337+, 567+, 838+
Pycnolejeunea dussiana, *470*
Pyke, David A. *See* Ponzetti, Jeanne M., 706+
Pyramidula tetragona, 109+
Pyramiduloideae, 112+
Pyrrhobryum bifarium, 538, *555*; dozyanum, 538; medium, 538; mnioides, 538, *555*; novae-caledoniae, 538; paramattense, 538; spiniforme, 538; vallis-gratiae, 538, *555*
Qiu, Yin-Long. *See* Hendry, Tory A., 738+
Quadrant production (*Cavicularia*), 457+*
Quandt, Dietmar. *See* Bell, Neil E., 533+
Quercus pyrenaica (lichens), 725
Racomitrioid leaf areolation, 96+
Racomitrium vulcanicola, 93
Racopilum cuspidigerum, 538; spectabile, 538
Radula, 746+
Ramalina farinacea, 730; fraxinea, 730; intermedia, 79+; roesleri, 529
Rank-free systematic treatment (Calymperaceae), 46+
*rbc*L (liverworts), 740
Reactive oxygen species, 101+
Recent literature on bryophytes—110(1), 133+; 110(2), 337+; 110(3), 567+; 110(4), 838+
Recent literature on lichens—204, 146+; —205, 351; —206, 577+; —207, 824+
Redwood epiphytes (lichens and bryophytes), 420+
Rehydration (bryophytes), 607+
Rehydrins (desiccation-tolerance), 607
Reiner-Drehwald, M. Elena and Anna Luiza Ilkiu-Borges. *Lejeunea huctumalcensis*, a widely distributed Lejeuneaceae from the Neotropics, and its relation to *Ceratolejeunea*, 465+
Renzaglia, Karen S., Scott Schutte, R. Joel Duff, Roberto Ligrone, A. Jonathan Shaw, Brent D. Mishler and Jeffrey G. Duckett. Bryophyte phylogeny: Advancing the molecular and morphological frontiers, 179+
Renzaglia, Karen S. *See* R. Joel Duff, 214+
Repeat-measurement error, 712
Reviewers of manuscripts submitted during 2006, 361+
Rhacocarpus purpurascens, 538
Rhacopilopsis trinitensis, 41
Rhizocarpon cinereovirens, 79+
Rhizogonium distichum, 538; graeffeanum, 538; novae-hollandiae, 538; pennatum, 538
Rhizomnium gracile, 538
Rhodobryum giganteum, 538
Rhytidiadelphus triquetrus, 538
Ribosomal DNA (*Nephroma*), 244+
Riccia, 746+
Rivas Plata, Eimy. *See* Lücking, Robert, 622+
RNA editing (hornworts), 220
Rock moss (harvest), 753+
ROS, 101+
Ros, Rosa M. *See* Werner, Olaf, 108+
Rosentreter, Roger. *See* DeBolt, Ann M., 254+
*rps*4 (liverworts), 740
Rutenbergia madagassa, 538
Salamanders (moss harvest), 761
Salazar Allen, Noris. *See* Villarreal, Juan Carlos, 279+
Salazinic acid (*Heterodermia*), 492
Santa Monica Mountains (*Lepraria*), 115+
SAS program (*Neckera*), 318
Scale (desiccation-tolerance), 597+
Scapania, 746+
Schistochila, 746+
Schofield, Wilfred B. *See* Talbot, Stephen S., 74+

Schultz, Matthias. On the identity of *Anema dodgei*, *Psorotichia segregata* and *Psorotichia squamulosa*, three misunderstood cyanolichens from the southwestern United States, 286+
Schumacher, René and Jiri Váňa. 2005. Identification keys to the liverworts and hornworts of Europe and Macaronesia (distribution and status) [review], 335+
Schuster, Rudolf M. *See* Hendry, Tory A., 738+
Schutte, Scott. *See* Renzaglia, Karen S., 179+
Sciadocladus, 556; celebensis, 556; kerrii, 556; menziesii, 556; **menziesii ssp. splendidum comb. nov.**, 556
Seasonal variation (desiccation-tolerance), 602+
Sequence editing and alignment (hornworts), 220
Sequencing (pleurocarps), 539
Sequoia epiphytes (lichens and bryophytes), 420+
Shaw, A. Jonathan. *See* Renzaglia, Karen S., 179+
Shaw, Blanka. *See* Háiková, Petra, 776+
Shaw, Jonathan (organizer). A molecular phylogenetic approach to the evolution of bryophytes, 178+
Shevock, James R. [Review of:] Bednarek-Ochyra, Halina. A taxonomic monograph of the moss genus *Codriophorus* P. Beauv. (Grimmiaceae), 130+
Shevock, James R. *See* Brinda, John C., 673+
Shoot increment (*Neckera*), 314+
Sillett, Stephen C. *See* Williams, Cameron B., 420+
SO_2, 724+
Soil chemistry (biotic soil crusts), 714
Soil-dwelling lichen and moss community composition, 707+
Sonesson, Mats, Bjartmar Sveinbjörnsson, Anders Tehler and Bengt Å. Carlsson. A comparison of the physiology, anatomy, and ribosomal DNA in alpine and subalpine populations of the lichen *Nephroma arcticum*—the effects of an eight-year transplant experiment, 244+
Sørensen distance measure, 712, 756
Spain (biomonitoring), 723+
Spatial distribution (*Bryopteris filicina*), 766+
Species area curves, 756
Species directly impacted by commercial moss harvest (list), 757+
Species diversity (lichens), 525+; (lichens and bryophytes), 429+
Species response curves, 778+
Species richness (biotic soil crusts), 713+; (lichens), 525+
Spence, John R. *See* Brinda, John C., 673+
Sperm cell evolution, front cover 110(2)*
Sphaerocarpos, 746+; texanus, 219+
Sphaerophorus globosus, 529
Sphaerosporoceros cf. adscendens, 218*+
Sphagnum (pores), 123;
Sphagnum section Cuspidata, 123; section Squarrosa, 123; section Subsecunda, 122; **atlanticum sp. nov.**, 274+*m; contortum, 123; fallax, 219+; **oregonense sp. nov.**, 122+*; palustre, 219+; subsecundum, 122; torreyanum, 274+*m
Spiridens camusii, 538; reinwardtii, 538; vieillardii, 538
Spore isthmus (*Caloplaca*), 798+
Spores (*Nothoceros*), 282*
Sporophyte development (desiccation-tolerance), 608+
Sporophyte evolution, 182+
Spring water chemistry, 779
Springs (Europe), 776+
Squamidium brasiliense, 24; macrocarpum, 42; nigricans, 10
Stark, Lloyd R. *See* Brinda, John C., 673+; *see* Proctor, Michael C. F., 595+
Steerecleus serrulatus, 759
Stem elongation (*Bazzania*), 304+
Stem water content (*Bazzania*), 301+
Steppe communities, 89
Sticta canariensis (distribution), 127
Stictic acid (*Usnea flammea*), 126
Stilbomyces berenice, 645
Stotler, Raymond E. *See* Freire, A. Virginia, 817+
Studlar, Susan Moyle and JeriLynn E. Peck. Commercial moss harvest in the Appalachian Mountains of West Virginia: targeted species and incidental take, 752+
Subfamilies (Funariaceae), 112+
Substrate effects (lichens and bryophytes), 437+
Subtropical forests (lichens), 254+
Succession (lichens and bryophytes), 438+
SURFER, 727
Sveinbjörnsson, Bjartman. *See* Sonesson, Mats, 244+
Symphyogyna, 746+
Symposium on a molecular phylogenetic approach to the evolution of bryophytes, 178+
Syrrhopodon (phylogenetic analysis), 46+
Syrrhopodon albo-vaginatus, 50; armatus, 50; ciliatus, 50; confertus, 50; croceus, 50; gardneri, 51+; incompletus, 51; involutus, 51; japonicus, 51; leprieurii, 51; loreus, 51; mahensis, 51; mauritianus, 51; muelleri, 51; parasiticus, 51; perarmatus, 51; prolifer, 51; tortilis, 51; trachyphyllus, 51; tristichus, 51
Systematic index to the bryophyte genera of Nevada, 704+
Takakia (phylogeny), 189*+
Takakia lepidozioides, 219+
Talbot, Stephen S., John W. Thomson and Wilfred B. Schofield. Lichens from Tetlin National Wildlife Refuge and vicinity, east-central Alaska, 74+
Targeted species (moss harvest), 752+
Targionia, 746+
Taxonomic classifications of Anthocerotophyta, 216*+
Taxonomy (pleurocarps), 533+
TBARS, 101+
Tehler, Anders. *See* Sonesson, Mats, 244+
Teloschistaceae, 798
Temnoma, 746+
Terpenoids (*Anaptychia bryorum*), 789; (*Anaptychia elbursiana*), 792; (*Anaptychia ulotrichoides*), 795
Terrain (pollutant distribution), 735
Terrón, A. *See* Fernández-Salegui, A. B., 723+
Tetlin National Wildlife Refuge (lichens), 74+*m
Tetraplodon mnioides, 538
Thalloid liverworts (phylogeny), 746+
Thamnobryum alopecurum, 539
Thiobarbituric acid, 101+
Thomson, John W. *See* Talbot, Stephen S., 74+
Thuidium delicatulum, 756+
Tønsberg, Tor. *See* Printzen, Christian, 487+
Topography (biotic soil crusts), 706+
Toren, David. *See* Yip, Kwok Leung, 510+
Touwiodendron gen. nov., 556; **diversifolium comb. nov.**, 557
Toxisensitive lichens, 733+
Toxitolerant lichens, 733+
Tracheophytes (desiccation-tolerance), 613*
Trachylejeunea (*Lejeunea*), 465
Trachyloma arborescens, 556; kerrii, 556
Transplant experiment (*Nephroma*), 244+
Trebouxia erici, 100+
Tree height, 771+
Tree of desiccation-tolerance in bryophytes, 174*
Tree orientation, 771+
Tree structure (lichens and bryophytes), 422+
Tree trunks (*Bryopteris filicina*), 766+
Tree-bisection-reconnection, 53
TreeRot.v2, 53
Treubia (phylogeny), 192*
Treubia lacunosa, 219+, 746+; pygmaea, 746+
Triandrophyllum, 746+
Tricharia, 623+; amazonum, 666; cuneata, 665*; aff. farinosa, 666; **floridensis sp. nov.**, 666*; helminthospora, 670; hyalina, 670; melanothrix, 668; planicarpa, 631; santesso-

nii, 667*; **subumbrosa sp. nov.**, 666+*m; **tuckerae sp. nov.**, 668+*; umbrosa, 668; vainioi, 666+*
Trichocolea, 746+
Trichotemnoma, 746+
Tripp, Erin A. *See* Lendemer, James C., 490+
trnL-trnF region (*Goniomitrium*), 109+
Tropical rain forest (disturbance), 767+
Tukey post-hoc tests, 778+
Tylimanthus, 746+
Typifications (*Fossombronia*), 817+
Ulota crispa, 539
United States, southeastern (*Bacidia*), 487+
United States, western (*Candelariella*), 365+, 375+, 391+
University of Porto herbarium, 119
Unweighted parsimony, 52
Uptake (*Trebouxia erici*), 100+
Usnea cornuta, 127; dalmatica, *126*; filipendula, 730; flammea, 126+m; fragilescens, 127; fulvoreagens, 127; halei, 127; longissima, 529; rubicunda, 127; rupestris, *126*; spp., 529; subfloridana, 127
Usneae (oceanic distribution), 127
Usnic acid (*Lecanora*), 485
Váczi, Peter. *See* Bačkor, Martin, 100+
Van Den Boom, P. P. G. *Lecanora crassithallina* (Lecanoraceae), a new lichen from northern Mexico, 482+
Variolaric acid (*Anaptychia bryorum*), 789+; (*Anaptychia isidiata*), 797; (*Anaptychia palmulata*), 793
Vegetation zones (Nevada), 675m
Vegetative desiccation-tolerance, 163+
Vellak, Kai. *See* Ingerpuu, Nele, 309+
Verrucaria aurella, *398*; cordeiroi, *120*
Vertical distribution (*Bryopteris filicina*), 768+
Vetiplanaxis pyrrhobryoides gen. et sp. nov., 518+*
Villarreal, Juan Carlos, Gabriela Hässel De Menéndez and Noris Salazar Allen. *Nothoceros superbus* (Dendrocerotaceae), a new hornwort from Costa Rica, 279+
Villarreal, Juan Carlos. *See* R. Joel Duff, 214+
Virtual bryophyte herbarium (New York Botanical Garden), 676
Vulpinic acid (*Candelariella*), 366+, 376+, 392+
Wall, Dennis P. *See* Fisher, Kirsten M., 46+
Wang, Bin. *See* Hendry, Tory A., 738+
Washington (biotic soil crusts), 706+
Water availability (*Bazzania*), 295+
Water conductivity, 778
Water relations (bryophytes), 599
Watson, Sereno, 674
Weather index (*Neckera*), 314+
Wedin, Mats. [Review of:] Holien, Håkon and Tor Tønsberg. Norsk lavflora, 129+
Werner, Olaf, Rosa M. Ros and Bernard Goffinet. A reconsideration of the systematic position of *Goniomitrium* (Funariaceae) based on chloroplast sequence markers, 108+
West Carpathians (*Philonotis*), 778
West Virginia (moss harvest), 752+
Westberg, Martin. *Candelariella* (Candelariaceae) in western United States and northern Mexico: 8-spored, lecanorine species, 391+; the polysporous species, 375+; the species with biatorine apothecia, 365+
Wetland habitats (*Philonotis*), 777+
Wetmore, Clifford M. Notes on *Caloplaca cerina* (Teloschistaceae) in North and Central America, 798+
Weymouthia cochlearifolia, 41
Williams, Cameron B. and Stephen C. Sillett. Epiphyte communities on redwood (*Sequoia sempervirens*) in northwestern California, 420+
Will-Wolf, Susan. [Review of:] Hill, David J. 2006. Surveying and report writing for lichenologists, 564+
Wilson, Paul. *See* Yip, Kwok Leung, 510+
Winds (pollutant distribution), 735
Wood, Andrew J. The nature and distribution of vegetative desiccation-tolerance in hornworts, liverworts, and mosses, 163+
Wood, Andrew J. *See* Proctor, Michael C. F., 595+
Xanthoria candelaria, 733+; parietina, 733+
Xanthorion community, 727+
Xenothallus, 746+
Xeric substrates (moss desiccation-tolerance), 599
Yahr, Rebecca. [Review of:] Archer, Alan. 2006. The lichen family Graphidaceae in Australia, 561+
Yang, Ya. *See* Hendry, Tory A., 738+
Yip, Kwok Leung, Paul Wilson, David Toren, and Ken M. Kellman. *Pleuridium mexicanum* (Ditrichaceae) new to the U.S.A. from California, 510+
Yip, Kwok Leung. *See* Fisher, Kirsten M., 46+
York, Peter V. *See* Bell, Neil E., 514+
Zeorin (*Heterodermia*), 492
Zoopsis, 499

THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
WILLIAM R. BUCK

Volume 110
2007

Contents

Allen, Bruce. See Buck et al.

Allen, Bruce and Robert E. Magill. A revision of *Orthostichella* (Neckeraceae) 1

Allen, Bruce and Robert R. Ireland. A new species of *Bartramia* (Bartramiaceae) from Chile 506

Alpert, Peter. See Proctor et al.

Anderson, Lewis Edward. [Death notice] 364

Andrus, Richard E. *Sphagnum atlanticum*, a new species from eastern North America 274

Andrus, Richard E. *Sphagnum oregonense* sp. nov. from western North America 123

Bačkor, Martin, Peter Váczi, Miloš Barták, Jana Buďová and Alexander Dzubaj. Uptake, photosynthetic characteristics and membrance lipid peroxidation levels in the lichen photobiont *Trebouxia erici* exposed to copper and cadmium 100

Barreno, E. See Fernández-Salegui et al.

Barták, Miloš. See Bačkor et al.

Bartholomew-Began, Sharon. See Jones and Bartholomew-Began.

Bednarek-Ochyra, Halina. See Köckinger et al.

Bell, Neil E., Dietmar Quandt, Terry J. O'Brien and Angela E. Newton. Taxonomy and phylogeny in the earliest diverging pleurocarps: square holes and bifurcating pegs 533

Bell, Neil E. and Peter V. York. *Vetiplanaxis pyrrhobryoides*, a new fossil moss genus and species from Middle Cretaceous Burmese amber 514

Braggins, John E. See Hendry et al.

Brinda, John C., Lloyd R. Stark, James R. Shevock and John R. Spence. An annotated checklist of the bryophytes of Nevada, with notes on collecting history in the state 673

Brunet, Jörg. See Caldiz et al.

Buck, William R. See Lücking et al.

Buck, William R., Bruce Allen and Ronald A. Pursell. Recent literature on bryophytes—110(1) 133

Buck, William R., Bruce Allen and Ronald A. Pursell. Recent literature on bryophytes—110(2) 337

Buck, William R., Bruce Allen and Ronald A. Pursell. Recent literature on bryophytes—110(3) 567

Buck, William R., Bruce Allen and Ronald A. Pursell. Recent literature on bryophytes—110(4) 838

Buďová, Jana. See Bačkor et al.

Caldiz, Mayra S., Jörg Brunet and Bengt Nihlgård. Lichen litter decomposition in *Nothofagus* forests of northern Patagonia: biomass and chemical changes over time 266

Cano, María J. See Jiménez and Cano.

Cargill, D. Christine. See Duff et al.

Carlsson, Bengt Å. See Sonesson et al.

Cleavitt, Natalie L., Karen L. Klima and Timothy J. Fahey. The role of water availability in seasonal growth in *Bazzania trilobata* on boulders in an eastern hemlock stand 295

Cleavitt, Natalie L. See Proctor et al.

Clerc, Philippe and Philip F. May. *Usnea flammea* (Lecanorales) new for North America 126

Davis, E. Christine. See Hendry et al.

DeBolt, Ann M., Roger Rosentreter and Erin P. Martin. Macrolichen diversity in subtropical forests of north-central Florida 254

Derr, Chiska C., Bruce McCune and Linda H. Geiser. Epiphytic macrolichen communities in *Pinus contorta* peatlands in southeastern Alaska 521

Duckett, Jeffrey G. See Renzaglia et al.

Duff, R. Joel. See Renzaglia et al.

Duff, R. Joel, Juan Carlos Villarreal, D. Christine Cargill and Karen S. Renzaglia. Progress and challenges toward developing a phylogeny and classification of the hornworts 214

Dzubaj, Alexander. See Bačkor et al.

Egan, Robert S. Recent literature on lichens—204 146

Egan, Robert S. Recent literature on lichens—205 351

Egan, Robert S. Recent literature on lichens—206 577

Egan, Robert S. Recent literature on lichens—207 824

Elix, John A. See Knudsen and Elix.

Equihua, Clementina and Miguel Equihua. Spatial distribution of *Bryopteris filicina* on tree trunks in the Selva Lacandona, Chiapas, Mexico 766

Equihua, Miguel. See Equihua and Equihua.

ESSLINGER, THEODORE L. A synopsis of the North American species of *Anaptychia* (Physciaceae) 788
FAHEY, TIMOTHY J. See CLEAVITT ET AL.
FERNÁNDEZ-SALEGUI, A. B., A. TERRÓN, E. BARRENO AND P. L. NIMIS. Biomonitoring with cryptogams near the power station of La Robla (León, Spain) 723
FISHER, KIRSTEN M., DENNIS P. WALL, KWOK LEUNG YIP AND BRENT D. MISHLER. Phylogeny of the Calymperaceae with a rank-free systematic treatment 46
FREIRE, A. VIRGINIA AND RAYMOND E. STOTLER. Typifications of the names of some Latin American species of *Fossombronia* (Fossombroniaceae) 817
GEISER, LINDA H. See DERR ET AL.
GOFFINET, BERNARD. See WERNER ET AL.
GOWARD, TREVOR AND BRUCE MCCUNE. *Hypogymnia canadensis* (Parmeliaceae), a new lichen from the Pacific Coast of North America 808
GREVEN, HENK. See HASTINGS AND GREVEN.
HÁJAK, MICHAL. See HÁJKOVÁ ET AL.
HÁJKOVÁ, PETRA, BLANKA SHAW, MICHAL HÁJAK, DIRK HINTERLAND AND VÍTĚZSLAV PLÁŠEK. The role of base saturation and altitude in habitat differentiation within *Philonotis* in springs and mires of three different European regions 776
HARRIS, RICHARD C. See LENDEMER ET AL.
HÄSSEL DE MENÉNDEZ, GABRIELA. See VILLARREAL ET AL.
HASTINGS, ROXANNE I. AND HENK GREVEN. *Grimmia milleri* sp. nov. (Grimmiaceae) from northeastern North America and the status of *Grimmia afroincurva* 500
HEDENÄS, LARS. Global diversity patterns among pleurocarpous mosses 319
HENDRY, TORY A., BIN WANG, YA YANG, E. CHRISTINE DAVIS, JOHN E. BRAGGINS, RUDOLF M. SCHUSTER AND YIN-LONG QIU. Evaluating phylogenetic positions of four liverworts from New Zealand, *Neogrollea notabilis*, *Jackiella curvata*, *Goebelobryum unguiculatum* and *Herzogianthus vaginatus*, using three chloroplast genes 738
HINTERLAND, DIRK. See HÁJKOVÁ ET AL.
ILKIU-BORGES, ANNA LUIZA. See REINER-DREHWALD AND ILKIU-BORGES.
INGERPUU, NELE, KAI VELLAK AND TÕNU MÕLS. Growth of *Neckera pennata*, an epiphytic moss of old-growth forests 309
IRELAND, ROBERT R. See ALLEN AND IRELAND.
JIMÉNEZ, JUAN A. AND MARÍA J. CANO. *Didymodon coquimbensis* (Pottiaceae), a new species from Chile 812
JONES, NICOLE L. AND SHARON BARTHOLOMEW-BEGAN. Juvenile gametophyte development in the Blasiales (Metzgeriidae). 2. Gemma/gemmaling ontogeny of *Cavicularia densa* 453
JOVAN, SARAH. [Review] Lichens and nitrogen 333
KELLMAN, KEN M. See YIP ET AL.
KLIMA, KAREN L. See CLEAVITT ET AL.
KNUDSEN, KERRY AND JOHN A. ELIX. A new *Lepraria* (Stereocaulaceae) from the Santa Monica Mountains in southern California 115
KNUDSEN, KERRY. [REVIEW] A crusty Pacific Northwest exposed 563
KÖCKINGER, HERIBERT, HALINA BEDNAREK-OCHYRA AND RYSZARD OCHYRA. *Bucklandiella nivalis* (Grimmiaceae), a new moss from the Alps of Austria 92
KONDRATYUK, SERGEY YA. In memoriam: Maria F. Makarevych, Dr. Sc., Professor (December 4, 1906–February 24, 1989) 480
KUČERA, JAN. [Review] New hepatic keys for Europe 335
LANE, DAVID M. See POWOROZNEK AND LANE.
LENDEMER, JAMES C., RICHARD C. HARRIS AND ERIN A. TRIPP. *Heterodermia neglecta* (Physciaceae), a new lichen species from eastern North America 490
LIGRONE, ROBERTO. See RENZAGLIA ET AL.
LÓPEZ DE SILANES, Mª EUGENIA. See PAZ-BERMÚDEZ AND LÓPEZ DE SILANES.
LÜCKING, ROBERT. The fungi *Microstelium hyalinum* and *Acrospermum puiggarii* are the same as the lichen *Gomphillus ophiosporus* (Ostropales: Gomphillaceae) 475
LÜCKING, ROBERT, WILLIAM R. BUCK AND EIMY RIVAS PLATA. The lichen family Gomphillaceae (Ostropales) in eastern North America, with notes on hyphophore development in *Gomphillus* and *Gyalideopsis* 622
MAGILL, ROBERT E. See ALLEN AND MAGILL
MANGOLD, ARMIN. [Review] New contribution to the systematics of the Thelotremataceae 332
MARTIN, ERIN P. See DEBOLT ET AL.

MAY, PHILIP F. See CLERC AND MAY.
MCCUNE, BRUCE. See DERR ET AL.
MCCUNE, BRUCE. See GOWARD AND MCCUNE.
MCCUNE, BRUCE. See PONZETTI ET AL.
MISHLER, BRENT D. See FISHER ET AL.
MISHLER, BRENT D. See PROCTOR ET AL.
MISHLER, BRENT D. See RENZAGLIA ET AL.
MÖLS, TÖNU. See INGERPUU ET AL.
MÜLLER, FRANK. *Meinungeria mouensis* (Lepidoziaceae), a new genus and species from New Caledonia 494
NEWTON, ANGELA E. See BELL ET AL.
NIHLGÅRD, BENGT. See CALDIZ ET AL.
NIMIS, P. L. See FERNÁNDEZ-SALEGUI ET AL.
O'BRIEN, TERRY J. See BELL ET AL.
OCHYRA, RYSZARD. See KÖCKINGER ET AL.
OLIVER, MELVIN J. See PROCTOR ET AL.
PAZ-BERMÚDEZ, GRACIELA AND MÁ EUGENIA LÓPEZ DE SILANES. The identity of two lichen species described from Portugal by J. Harmand 119
PECK, JERILYNN E. See MOYLE AND PECK.
PLÁŠEK, VÍTĚZSLAV. See HÁJKOVÁ ET AL.
PONZETTI, JEANNE M., BRUCE MCCUNE AND DAVID A. PYKE. Biotic soil crusts in relation to topography, cheatgrass and fire in the Columbia Basin, Washington 706
POWOROZNEK, EMILY L. AND DAVID M. LANE. Index to volume 110 852
PRINTZEN, CHRISTIAN AND TOR TØNSBERG. *Bacidia lobarica* (Bacidiaceae, Lecanorales) sp. nov., a sorediate lichen from the southeastern U.S.A. 487
PROCTOR, MICHAEL C. F., MELVIN J. OLIVER, ANDREW J. WOOD, PETER ALPERT, LLOYD R. STARK, NATALIE L. CLEAVITT AND BRENT D. MISHLER. Desiccation-tolerance in bryophytes: a review 595
PURSELL, RONALD A. [Review] More sumptuous bryophyte photographs 566
PURSELL, RONALD A. See BUCK ET AL.
PYKE, DAVID A. See PONZETTI ET AL.
QIU, YIN-LONG. See HENDRY ET AL.
QUANDT, DIETMAR. See BELL ET AL.
REINER-DREHWALD, M. ELENA AND ANNA LUIZA ILKIU-BORGES. *Lejeunea huctumalcensis*, a widely distributed Lejeuneaceae from the Neotropics, and its relation to *Ceratolejeunea* 465
RENZAGLIA, KAREN S. See DUFF ET AL.
RENZAGLIA, KAREN S., SCOTT SCHUTTE, R. JOEL DUFF, ROBERTO LIGRONE, A. JONATHAN SHAW, BRENT D. MISHLER AND JEFFREY G. DUCKETT. Bryophyte phylogeny: advancing the molecular and morphological frontiers 179
RIVAS PLATA, EIMY. See LÜCKING ET AL.
ROS, ROSA M. See WERNER ET AL.
ROSENTRETER, ROGER. See DEBOLT ET AL.
SALAZAR ALLEN, NORIS. See VILLARREAL ET AL.
SCHOFIELD, WILFRED B. See TALBOT ET AL.
SCHULTZ, MATTHIAS. On the identity of *Anema dodgei*, *Psorotichia segregata* and *Psorotichia squamulosa*, three misunderstood cyanolichens from the southwestern United States 286
SCHUSTER, RUDOLF M. See HENDRY ET AL.
SCHUTTE, SCOTT. See RENZAGLIA ET AL.
SHAW, A. JONATHAN. See RENZAGLIA ET AL.
SHAW, BLANKA. See HÁJKOVÁ ET AL.
SHAW, JONATHAN. [Symposium] A molecular phylogenetic approach to the evolution of bryophytes 178
SHEVOCK, JAMES R. [Review] New monograph of a *Racomitrium* segregate 130
SHEVOCK, JAMES R. See BRINDA ET AL.
SILLETT, STEPHEN C. See WILLIAMS AND SILLETT.
SONESSON, MATS, BJARTMAR SVEINBJÖRNSSON, ANDERS TEHLER AND BENGT Å. CARLSSON. A comparison of the physiology, anatomy and ribosomal DNA in alpine and subalpine populations of the lichen *Nephroma arcticum*—the effects of an eight-year transplant experiment 244
SPENCE, JOHN R. See BRINDA ET AL.
STARK, LLOYD R. See BRINDA ET AL.

STARK, LLOYD R. See PROCTOR ET AL.
STOTLER, RAYMOND E. See FREIRE AND STOTLER.
STUDLAR, SUSAN MOYLE AND JERILYNN E. PECK. Commercial moss harvest in the Appalachian Mountains of West Virginia: targeted species and incidental take 752
SVEINBJÖRNSSON, BIARTMAR. See SONESSON ET AL.
TALBOT, STEPHEN S., JOHN W. THOMSON AND WILFRED B. SCHOFIELD. Lichens from Tetlin National Refuge and vicinity, east-central Alaska 74
TEHLER, ANDERS. See SONESSON ET AL.
TERRÓN, A. See FERNÁNDEZ-SALEGUI ET AL.
THOMSON, JOHN W. See TALBOT ET AL.
TØNSBERG, TOR. See PRINTZEN AND TØNSBERG.
TOREN, DAVID. See YIP ET AL.
TRIPP, ERIN A. See LENDEMER ET AL.
VÁCZI, PETER. See BAČKOR ET AL.
VAN DEN BOOM, P. P. G. *Lecanora crassithallina* (Lecanoraceae), a new lichen species from northern Mexico 482
VELLAK, KAI. See INGERPUU ET AL.
VILLARREAL, JUAN CARLOS. See DUFF ET AL.
VILLARREAL, JUAN CARLOS, GABRIELA HÄSSEL DE MENÉNDEZ AND NORIS SALAZAR ALLEN. *Nothoceros superbus* (Dendrocerotaceae), a new hornwort from Costa Rica 279
WALL, DENNIS P. See FISHER ET AL.
WANG, BIN. See HENDRY ET AL.
WEDIN, MATS. [Review] A new Norwegian lichen flora 129
WERNER, OLAF, ROSA M. ROS AND BERNARD GOFFINET. A reconsideration of the systematic position of *Goniomitrium* (Funariaceae) based on chloroplast sequence markers 108
WESTBERG, MARTIN. *Candelariella* (Candelariaceae) in western United States and northern Mexico: the 8-spored, lecanorine species 391
WESTBERG, MARTIN. *Candelariella* (Candelariaceae) in western United States and northern Mexico: the polysporous species 375
WESTBERG, MARTIN. *Candelariella* (Candelariaceae) in western United States and northern Mexico: the species with biatorine apothecia 365
WETMORE, CLIFFORD M. Notes on *Caloplaca cerina* (Teloschistaceae) in North and Central America 798
WILLIAMS, CAMERON B. AND STEPHEN C. SILLETT. Epiphyte communities on redwood (*Sequoia sempervirens*) in northwestern California 420
WILL-WOLF, SUSAN. [Review] Dotting the i's for contracted lichen (and bryophyte?) surveys 564
WILSON, PAUL. See YIP ET AL.
WOOD, ANDREW J. The nature and distribution of vegetative desiccation-tolerance in hornworts, liverworts and mosses 163
WOOD, ANDREW J. See PROCTOR ET AL.
YAHR, REBECCA. [Review] A new look at Australian graphids 561
YANG, YA. See HENDRY ET AL.
YIP, KWOK LEUNG. See FISHER ET AL.
YIP, KWOK LEUNG, PAUL WILSON, DAVID TOREN AND KEN M. KELLMAN. *Pleuridium mexicanum* (Ditrichaceae) new to the U.S.A. from California 510
YORK, PETER V. See BELL AND YORK.